

THE EQUIVARIANT TODA LATTICE, I

EZRA GETZLER

The Toda lattice is an infinite dimensional dynamical system of commuting flows

$$(\partial, \delta_n, \bar{\delta}_n \mid n > 0),$$

acting on functions $(q, a_k, \bar{a}_k \mid k > 0)$ defined on a one-dimensional lattice. In the limit of small lattice spacing ε , which is all that will concern us here (Takasaki and Takebe [10]), the functions $(q, a_k, \bar{a}_k)$ become functions of a real parameter x , and the role of translation by one unit of the lattice is taken by the operator $E = e^{\varepsilon \partial}$, where ∂ is the infinitesimal generator of translations in x .

The derivations δ_1 and $\bar{\delta}_1$ act on the variables q , a_1 and $\bar{a}_1$ by the formulas

$$(1) \quad \delta_1 \bar{a}_1 = \bar{\delta}_1 a_1 = \nabla q, \quad \delta_1 \log q = \nabla a_1, \quad \bar{\delta}_1 \log q = \nabla \bar{a}_1,$$

where $\nabla : \mathcal{A} \rightarrow \mathcal{A}$ is the infinite-order differential operator

$$\begin{aligned} \nabla &= \varepsilon^{-1} (E^{1/2} - E^{-1/2}) = \sum_{k=0}^{\infty} \frac{\varepsilon^{2k} \partial^{2k+1}}{2^{2k} (2k+1)!} \\ &= \partial + \frac{1}{24} \varepsilon^2 \partial^3 + O(\varepsilon^4). \end{aligned}$$

These formulas imply the **Toda equation**:

$$(2) \quad \delta_1 \bar{\delta}_1 \log q = \nabla^2 q.$$

The higher Toda flows are symmetries of this equation.

An abstract mathematical formulation of the Toda lattice is obtained by realizing the derivations $(\partial, \delta_n, \bar{\delta}_n)$ on the free differential algebra $\mathcal{A} = \mathbb{Q}_\varepsilon \{q, a_k, \bar{a}_k \mid k > 0\}$, defined over the ring $\mathbb{Q}_\varepsilon = \mathbb{Q}[[\varepsilon]]$. Reductions of the Toda lattice are differential ideals $\mathcal{I}$ in $\mathcal{A}$ invariant under conjugation and closed under the derivations δ_n and $\bar{\delta}_n$. For example, the **Toda chain** is described by the differential ideal with generators

$$\{a_1 - \bar{a}_1, a_2 - q, \bar{a}_2 - q, a_k, \bar{a}_k \mid k > 2\}.$$

In this paper, we study a reduction of the Toda lattice, which we call the equivariant Toda lattice. If ν is a formal parameter, this reduction is defined by the following constraint on the Lax operator:

$$(3) \quad (\delta_1 - \bar{\delta}_1)L = \nu \partial L.$$

We prove that the corresponding differential ideal $\mathcal{I}_\nu \subset \mathcal{A}[\nu]$ is isomorphic to the differential algebra

$$\mathbb{Q}_{\varepsilon,\nu}\{q, v, \bar{v}\}/(\nu\partial q - \nabla(v - \bar{v})) \otimes_{\mathbb{Q}_{\varepsilon,\nu}} \mathbb{Q}_{\varepsilon,\nu}[z_k, \bar{z}_k \mid k > 0],$$

where $\mathbb{Q}_{\varepsilon,\nu} = \mathbb{Q}_\varepsilon[\nu]$, v and $\bar{v}$ are the images of a_1 and $\bar{a}_1 \in \mathcal{A}$, and z_k and $\bar{z}_k$ are constants of motion, which may be defined by the following equation:

$$\left(L - \nu + \sum_{k=1}^{\infty} z_k L^{-k}\right) \frac{\partial L}{\partial v} = L.$$

The equivariant Toda lattice may be used to describe the equivariant Gromov-Witten invariants of $\mathbb{CP}^1$. Let $\mathbb{T}$ be the multiplicative group of $\mathbb{C}$, and let X be a topological space with an action of $\mathbb{T}$. The equivariant cohomology $H_{\mathbb{T}}^*(X, \mathbb{Z})$ of X is a module over the graded ring $H_{\mathbb{T}}^\bullet(*, \mathbb{Z}) \cong \mathbb{Z}[\nu]$, where $\nu \in H_{\mathbb{T}}^2(*, \mathbb{Z})$. The equivariant cohomology $H_{\mathbb{T}}^\bullet(\mathbb{CP}^1, \mathbb{Z})$ of the projective line $\mathbb{CP}^1$ admits a presentation

$$H_{\mathbb{T}}^\bullet(\mathbb{CP}^1, \mathbb{Z}) \cong \mathbb{Z}[H, \nu]/(H(H - \nu)),$$

where H is the equivariant Chern class $c_1(\mathcal{O}(1)) \in H_{\mathbb{T}}^2(\mathbb{CP}^1, \mathbb{Z})$.

Denote the k th descendants of the cohomology classes 1 and H in Gromov-Witten theory by $\tau_{k,P}$ and $\tau_{k,Q}$ respectively; also, abbreviate $\tau_{0,P}$ and $\tau_{0,Q}$ to P and Q . The genus 0 equivariant Gromov-Witten invariants of $\mathbb{CP}^1$ are integrals over the moduli space $\overline{\mathcal{M}}_{0,n}(\mathbb{CP}^1)$ of stable maps of genus g with n marked points:

$$\langle \tau_{k_1,P} \dots \tau_{k_m,P} \tau_{\ell_1,Q} \dots \tau_{\ell_n,Q} \rangle_g \in H_{\mathbb{T}}^\bullet(*, \mathbb{Q}) \cong \mathbb{Q}[\nu].$$

The large phase space is the formal affine space with coordinates $\{s_k, t_k \mid k \geq 0\}$. The genus g Gromov-Witten potential $\mathcal{F}_g$ of $\mathbb{CP}^1$ is the generating function on the large phase space given by the formula

$$\mathcal{F}_g = \sum_{m,n=0}^{\infty} \frac{1}{m!n!} \sum_{\substack{k_1, \dots, k_m \\ \ell_1, \dots, \ell_n}} s_{k_1} \dots s_{k_m} t_{\ell_1} \dots t_{\ell_n} \langle \tau_{k_1,P} \dots \tau_{k_m,P} \tau_{\ell_1,Q} \dots \tau_{\ell_n,Q} \rangle_g.$$

We may combine the Gromov-Witten potentials into a single generating function by interpreting ε as a genus expansion parameter, and writing

$$\mathcal{F} = \sum_{g=0}^{\infty} \varepsilon^{2g} \mathcal{F}_g.$$

Based on explicit calculations using the topological recursion relations in genus 0 and 1, Pandharipande conjectured [9] that the following equation holds for the total Gromov-Witten potential:

$$(4) \quad \partial_0 \bar{\partial}_0 \mathcal{F} = \exp(\nabla^2 \mathcal{F}).$$

Here, $\nabla = \varepsilon^{-1}(\mathbf{E}^{1/2} - \mathbf{E}^{-1/2})$, where $\mathbf{E} = e^{\varepsilon \partial}$, and $\partial = \partial/\partial s_0$. This equation was proved recently by Okounkov and Pandharipande [7].

On applying the operator ∇^2 to both sides of (4) and identifying the vector fields ∂_0 and $\bar{\partial}_0$ with the Toda flows δ_1 and $\bar{\delta}_1$, we obtain the Toda equation (2) for $q = \exp(\nabla^2 \mathcal{F})$. Observe that $\partial_1 - \bar{\partial}_1 = \nu \partial$; this equation is formally identical to the constraint defining the equivariant Toda lattice.

The equivariant Toda lattice and the equivariant Gromov-Witten theory of $\mathbb{CP}^1$ each involve sequences $\{\delta_n, \bar{\delta}_n\}$ and $\{\partial_n = \partial/\partial t_n, \bar{\partial}_n = \partial/\partial t_n - \nu \partial/\partial s_n\}$ of commuting derivations, in the first case on the algebra $\mathcal{C}$, and in the second case on functions on the large phase space. These sequences of vector fields may be compared by means of a morphism

$$\mathcal{A}[\nu]/\mathcal{I}_\nu \longrightarrow \mathbb{Q}_{\varepsilon, \nu} \llbracket s_k, t_k \mid k \geq 0 \rrbracket$$

of differential algebras which sends the generators q , v and $\bar{v}$ to $\exp(\nabla^2 \mathcal{F})$, $\nabla \partial_0 \mathcal{F}$ and $\nabla \bar{\partial}_0 \mathcal{F}$, and the constants z_k and $\bar{z}_k$ to 0. In fact, the following relationship between these flows holds:

$$(5a) \quad \sum_{k=0}^{\infty} z^{k+1} \partial_k = \sum_{n=1}^{\infty} \frac{z^n \delta_n}{(1+z\nu)(2+z\nu)\dots(n+z\nu)},$$

$$(5b) \quad \sum_{k=0}^{\infty} z^{k+1} \bar{\partial}_k = \sum_{n=1}^{\infty} \frac{z^n \bar{\delta}_n}{(1-z\nu)(2-z\nu)\dots(n-z\nu)}.$$

We conjectured this in a preprint of this paper, based on a proof of the result in genus 0 (see Section 5), together with calculations in genus 1 for small values of n ; it has recently been proved by Okounkov and Pandharipande [7]. Thus, the equivariant Toda lattice yields a description of the equivariant Gromov-Witten invariants of $\mathbb{CP}^1$ in terms of a Lax operator whose coefficients are obtained by an explicit recursion.

In a sequel to this paper, we relate the equivariant Toda lattice to the dressing operator formalism. Let $\log(L) = W \log(\Lambda) W^{-1}$ be the logarithm of the Lax operator L , related to the operator $\ell = \varepsilon(\partial W) W^{-1}$ by the formula

$$\log(L) = \log(\Lambda) - \ell.$$

Borrowing ideas of Carlet, Dubrovin and Zhang [1], we show that the equivariant Toda lattice may be characterized by the expansion

$$\Lambda + v + q\Lambda^{-1} = L + \nu\ell - \sum_{k=1}^{\infty} \frac{z_k}{k} L^{-k}.$$

In particular, the equation $(\delta_1 - \bar{\delta}_1)W = \nu \partial W$ is equivalent to the vanishing of the coefficients z_k . Under the same hypothesis, we also show that the equivariant Toda lattice is Hamiltonian; this gives a more direct relationship between the results of Okounkov and Pandharipande [7] and the original Toda conjecture (Eguchi and Yang [3], Eguchi, Hori and Yang [4], Pandharipande [8], Getzler [5]).

Acknowledgements. This paper has its origin in discussions with T. Eguchi and R. Pandharipande at the workshop on Duality in Mirror Symmetry at the Institute for Theoretical Physics of the University of California, Santa Barbara. It was written during a membership of the Institute for Advanced Study, Princeton. We are grateful to R. Dijkgraaf, B. Dubrovin and Y. Zhang for useful discussions.

The author is partially supported by the NSF under grant DMS-0072508, and, through the Institute for Advanced Study, under grant DMS-9729992.

APPENDIX A. ANOTHER FORMULATION OF THE EQUIVARIANT TODA CONJECTURE

In this appendix, we invert the relations of (5a) and (5b). Define the (unsigned) Stirling numbers (of the first kind) $\begin{bmatrix} n \\ k \end{bmatrix}$ by the generating function

$$\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} \nu^k = \prod_{j=0}^{n-1} (\nu + j).$$

Proposition A.1.

$$(6a) \quad \delta_n = n \sum_{k=1}^n \nu^{k-1} \begin{bmatrix} n \\ k \end{bmatrix} \partial_{n-k},$$

$$(6b) \quad \bar{\delta}_n = n \sum_{k=1}^n (-\nu)^{k-1} \begin{bmatrix} n \\ k \end{bmatrix} \bar{\partial}_{n-k}.$$

Proof. We prove the equivalence of (5a) with (6a); the equivalence of (5b) and (6b) is similar. Equation (5a) may be restated as saying that

$$\partial_k = \sum_{n=1}^{k+1} (-\nu)^{k-n+1} h_{k-n+1} \left(1, \frac{1}{2}, \dots, \frac{1}{n}\right) \frac{\delta_n}{n!},$$

where h_ℓ is the complete symmetric polynomial of degree ℓ . We wish to prove that

$$\frac{\delta_n}{n!} = \sum_{\ell=0}^{n-1} \nu^{n-\ell-1} e_{n-\ell-1} \left(1, \frac{1}{2}, \dots, \frac{1}{n-1}\right) \partial_\ell,$$

where e_ℓ is the elementary symmetric polynomial of degree ℓ . In other words, we wish to prove that

$$\sum_{\ell=0}^{n-1} \nu^{n-\ell-1} (-\nu)^{\ell-m+1} e_{n-\ell-1} \left(1, \frac{1}{2}, \dots, \frac{1}{n-1}\right) h_{\ell-m+1} \left(1, \frac{1}{2}, \dots, \frac{1}{m}\right) = \delta_{n,m}.$$

This is clearly true if $n \leq m$; thus, we have only to prove that the left-hand side vanishes when $n > m$. In this case, it equals ν^{n-m} times the coefficient of ν^{n-m} in the generating function

$$\prod_{j=1}^{n-1} (1 + j\nu) \cdot \prod_{j=1}^m (1 + j\nu)^{-1} = \prod_{j=m+1}^{n-1} (1 + j\nu),$$

which is a polynomial of degree $n - m - 1$; hence, the coefficient in question vanishes. $\square$

2. DIFFERENCE OPERATORS

In this section, we recall the mathematical structure underlying the Toda lattice; this material is adapted from the fundamental papers of Ueno and Takasaki [11] and Kupershmidt [6].

All of the commutative algebras which we consider in this paper carry an involution $p \mapsto \bar{p}$, and all ideals which we consider are closed under this involution. By a **differential algebra**, we mean a commutative algebra with derivation ∂ such that

$$\partial \bar{p} = \overline{\partial p}.$$

A **differential ideal** is an ideal closed under the action of the differential ∂ . If S is a subset of a differential algebra $\mathcal{A}$, denote the differential ideal generated by $S \cup \bar{S}$ by (S) , where $\bar{S} = \{\bar{x} \mid x \in S\}$ is the conjugate of S .

If $\mathcal{A}$ is a differential algebra and S is a set, the free differential algebra $\mathcal{A}\{S\}$ generated by S is the polynomial algebra

$$\mathcal{A}[\partial^n x, \partial^n \bar{x} \mid x \in S, n \geq 0],$$

with differential $\partial(\partial^n x) = \partial^{n+1} x$.

An **evolutionary derivation** δ of a differential algebra $\mathcal{A}$ is a derivation such that $[\partial, \delta] = 0$. The evolutionary derivations form a Lie subalgebra of the Lie algebra of derivations of $\mathcal{A}$, with involution

$$\bar{\delta} p = \overline{\delta \bar{p}}.$$

Let $\mathcal{A}$ be a differential algebra over $\mathbb{Q}_\varepsilon$, and let $q \in \mathcal{A}$ be a regular element (that is, having no zero-divisors) such that $\bar{q} = q$. The localization $q^{-1}\mathcal{A}$ of $\mathcal{A}$ is a filtered differential algebra, with differential $\partial(q^{-1}) = -q^{-2}\partial q$. Let $\Phi_\pm(\mathcal{A}, q)$ be the associative algebras of **difference operators**

$$\begin{aligned} \Phi_+(\mathcal{A}, q) &= \left\{ \sum_{k=-\infty}^{\infty} p_k \Lambda^k \mid p_k \in q^{-k}\mathcal{A}, p_k = 0 \text{ for } k \ll 0 \right\}, \\ \Phi_-(\mathcal{A}, q) &= \left\{ \sum_{k=-\infty}^{\infty} p_k \Lambda^k \mid p_k \in \mathcal{A}, p_k = 0 \text{ for } k \gg 0 \right\}, \end{aligned}$$

with product

$$\sum_i a_i \Lambda^i \cdot \sum_j b_j \Lambda^j = \sum_k \left(\sum_{i+j=k} (\mathbb{E}^{-j/2} a_i) (\mathbb{E}^{i/2} b_j) \right) \Lambda^k.$$

Note that $\Phi_-(\mathcal{A}, q)$ is in fact independent of q .

Let $A \mapsto A_{\pm}$ be the projections on $\Phi_{\pm}(\mathcal{A}, q)$ defined by the formulas

$$\left(\sum_{k=-\infty}^{\infty} p_k \Lambda^k \right)_+ = \sum_{k=0}^{\infty} p_k \Lambda^k, \quad \left(\sum_{k=-\infty}^{\infty} p_k \Lambda^k \right)_- = \sum_{k=-\infty}^{-1} p_k \Lambda^k.$$

We see that $A = A_- + A_+$. Define the residue $\text{res} : \Phi_p m(\mathcal{A}, q) \rightarrow \mathcal{A}$ by the formula

$$\text{res} \left(\sum_{k=-\infty}^{\infty} p_k \Lambda^k \right) = p_0.$$

For $k \in \mathbb{Z}$, let $[k]$ be the isomorphism of $\mathcal{A}$

$$[k] = \frac{\mathbf{E}^{k/2} - \mathbf{E}^{-k/2}}{\mathbf{E}^{1/2} - \mathbf{E}^{-1/2}} = \sum_{j=1}^k \mathbf{E}^{(k+1)/2-j} = k + O(\varepsilon^2).$$

Define $q^{[k]}$ by the recursion

$$q^{[k+1]} = \mathbf{E}^k q \cdot \mathbf{E}^{-1/2} q^{[k]},$$

with initial condition $q^{[0]} = 1$. The involution

$$A = \sum_{k=-\infty}^{\infty} p_k \Lambda^k \mapsto \bar{A} = \sum_{k=1}^{\infty} \bar{p}_k q^{[k]} \Lambda^{-k} + \bar{p}_0 + \sum_{k=1}^{\infty} \bar{p}_{-k} q^{-[k]} \Lambda^k,$$

defines an anti-isomorphism between the algebras $\Phi_+(\mathcal{A}, q)$ and $\Phi_-(\mathcal{A}, q)$.

3. THE TODA LATTICE

To formulate the Toda lattice, we introduce the differential algebra

$$\mathcal{A} = \mathbb{Q}_{\varepsilon} \{q, a_k \mid k > 0\} / (q - \bar{q}).$$

Since the generator a_1 plays a special role in the theory, we denote it by v . It will be useful to define the symbol a_0 to equal 1.

The Lax operator of the Toda lattice is the difference operator

$$L = \Lambda + \sum_{k=1}^{\infty} a_k \Lambda^{-k+1} \in \Phi_-(\mathcal{A}, q);$$

its conjugate $\bar{L}$ is given by the formula

$$\bar{L} = q \Lambda^{-1} + \sum_{k=1}^{\infty} \bar{a}_k q^{-[k-1]} \Lambda^{k-1} \in \Phi_+(\mathcal{A}, q).$$

Introduce elements $p_k(n) \in \mathcal{A}$, defined for all $n \geq 0$ and $k \in \mathbb{Z}$:

$$L^n = \sum_{k=-\infty}^n p_k(n) \Lambda^k.$$

To define the evolutionary derivation δ_n on the generators a_k of $\mathcal{A}$, introduce the difference operator $B_n = L_+^n$, and impose the Lax equation $\delta_n L = \varepsilon^{-1}[B_n, L]$. This equation means that

$$\varepsilon^{-1}[B_n, L] = \sum_{k=1}^{\infty} \delta_n a_k \Lambda^{-k+1}.$$

In order for this to be meaningful, it must be shown that the coefficient of Λ^k in $[B_n, L]$ vanishes for $k > 0$. This follows from the identity $[L^n, L] = 0$: we have

$$[B_n, L] = [B_n, L] - [L^n, L] = -[L_-^n, L],$$

and it is clear that the coefficient of Λ^k in $[L_-^n, L]$ vanishes if $k > 0$. We also see that $\delta_n a_k$ equals the coefficient of Λ^{-k+1} in

$$-\varepsilon^{-1} \sum_{j=0}^k [p_{j-k}(n) \Lambda^{j-k}, a_j \Lambda^{-j+1}],$$

hence that

$$\begin{aligned} (7) \quad \delta_n a_k &= \nabla p_{-k}(n) + \varepsilon^{-1} \sum_{j=1}^{k-1} \left(\mathbb{E}^{(1-j)/2} p_{j-k}(n) \mathbb{E}^{(k-j)/2} a_j - \mathbb{E}^{(j-1)/2} p_{j-k}(n) \mathbb{E}^{(j-k)/2} a_j \right) \\ &= \nabla p_{-k}(n) + \sum_{j=1}^{k-1} \left(\mathbb{E}^{(1-j)/2} p_{j-k}(n) \nabla[k-j] a_j - \mathbb{E}^{(j-k)/2} a_j \nabla[j-1] p_{j-k}(n) \right). \end{aligned}$$

In particular, $\delta_n v = \nabla p_{-1}(n)$.

To define δ_n on the remaining generators q and $\bar{a}_k$ of $\mathcal{A}$, we impose the Lax equation $\delta_n \bar{L} = \varepsilon^{-1}[B_n, \bar{L}]$. In particular, we see that

$$(8) \quad \delta_n q = q \nabla p_0(n),$$

and hence that $\delta_n q^{[k]} = q^{[k]} \nabla[k] p_0(n)$. It also follows that

$$\delta_n (q^{-[k-1]} \bar{a}_k) = q^{-[k-1]} (\delta_n \bar{a}_k - \bar{a}_k \nabla[k-1] p_0(n))$$

equals the coefficient of Λ^{k-1} in

$$\varepsilon^{-1} \sum_{j=0}^k [p_{k-j}(n) \Lambda^{k-j}, q^{-[j-1]} \bar{a}_j \Lambda^{j-1}],$$

hence that

(9)

$$\begin{aligned}
\delta_n \bar{a}_k &= \bar{a}_k \nabla[k-1]p_0(n) \\
&\quad + \varepsilon^{-1} q^{[k-1]} \sum_{j=0}^k \left(\mathbb{E}^{(1-j)/2} p_{k-j}(n) \mathbb{E}^{(k-j)/2} q^{-[j-1]} \bar{a}_j - \mathbb{E}^{(j-1)/2} p_{k-j}(n) \mathbb{E}^{(j-k)/2} q^{-[j-1]} \bar{a}_j \right) \\
&= \bar{a}_k \nabla[k-1]p_0(n) \\
&\quad + \varepsilon^{-1} \sum_{j=0}^k \left(\mathbb{E}^{(1-j)/2} (q^{[k-j]} p_{k-j}(n)) \mathbb{E}^{(k-j)/2} \bar{a}_j - \mathbb{E}^{(j-1)/2} (q^{[k-j]} p_{k-j}(n)) \mathbb{E}^{(j-k)/2} \bar{a}_j \right) \\
&= \nabla(q^{[k]} p_k(n)) \\
&\quad + \sum_{j=1}^{k-1} \left(\mathbb{E}^{(1-j)/2} (q^{[k-j]} p_{k-j}(n)) \nabla[k-j] \bar{a}_j - \mathbb{E}^{(j-k)/2} \bar{a}_j \nabla[j-1] (q^{[k-j]} p_{k-j}(n)) \right).
\end{aligned}$$

In particular, $\delta_n \bar{v} = \nabla(qp_1(n))$.

Denote by $\alpha : \mathcal{A} \rightarrow \mathbb{Q}_\varepsilon$ the homomorphism which sends the generators $\{q, a_k, \bar{a}_k\}$ of $\mathcal{A}$ to 0. By formulas (7), (8) and (9), we see that $\delta_n q$, $\delta_n a_k$ and $\delta_n \bar{a}_k$ all lie in the ideal $(\partial \mathcal{A})$ of $\mathcal{A}$, and hence

$$(10) \quad \alpha \cdot \delta_n = 0.$$

We now recall the proof that the derivations δ_m and δ_n commute. The proof relies on the Zakharov-Shabat equation

$$(11) \quad \delta_m B_n - \delta_n B_m = \varepsilon^{-1} [B_m, B_n].$$

To prove this equation, observe that

$$\begin{aligned}
\delta_m B_n &= (\delta_m L^n)_+ = \varepsilon^{-1} [B_m, L^n]_+ = \varepsilon^{-1} [B_m, B_n + L_-^n]_+ \\
&= \varepsilon^{-1} [B_m, B_n] + \varepsilon^{-1} [B_m, L_-^n]_+.
\end{aligned}$$

Since $[L^m, L^n] = 0$, we also see that

$$[B_m, L^n]_+ = [L^m - L_-^m, L^n]_+ = -[L_-^m, B_n]_+.$$

It follows that

$$\delta_m B_n - \delta_n B_m = \varepsilon^{-1} ([B_m, B_n] + [B_m, L_-^n]_+) + \varepsilon^{-1} [L_-^n, B_m]_+ = \varepsilon^{-1} [B_m, B_n].$$

From (11), we easily see that the derivations δ_m and δ_n commute:

$$\begin{aligned}
[\delta_m, \delta_n] L &= \varepsilon^{-1} \delta_m [B_n, L] - \varepsilon^{-1} \delta_n [B_m, L] \\
&= \varepsilon^{-1} [\delta_m B_n, L] + \varepsilon^{-2} [B_n, [B_m, L]] - \varepsilon^{-1} [\delta_n B_m, L] - \varepsilon^{-2} [B_m, [B_n, L]] \\
&= \varepsilon^{-1} [\delta_m B_n - \delta_n B_m, L] + \varepsilon^{-2} [B_n, [B_m, L]] - \varepsilon^{-2} [B_m, [B_n, L]] = 0.
\end{aligned}$$

The derivation $\bar{\delta}_n$ is defined to be the conjugate of δ_n , acting on the generators of $\mathcal{A}$ by the formulas

$$\bar{\delta}_n q = \overline{\delta_n q}, \quad \bar{\delta}_n a_k = \overline{\delta_n a_k}, \quad \bar{\delta}_n \bar{a}_k = \overline{\delta_n a_k}.$$

The following proposition establishes the Lax equation for this derivation.

Proposition 3.1. *Let $C_n = -\bar{L}^n$; then $\bar{\delta}_n L = \varepsilon^{-1}[C_n, L]$ and $\bar{\delta}_n \bar{L} = \varepsilon^{-1}[C_n, \bar{L}]$.*

Proof. We have

$$\begin{aligned} \overline{\bar{\delta}_n L} &= \overline{\bar{\delta}_n q} \Lambda^{-1} + \sum_{k=1}^{\infty} \overline{q^{-(k-1)} (\bar{\delta}_n \bar{a}_k - \bar{a}_k q^{-(k-1)} \bar{\delta}_n q^{(k-1)})} \Lambda^{k-1} \\ &= \sum_{k=1}^{\infty} \overline{\bar{\delta}_n \bar{a}_k} \Lambda^{-k+1} + \nabla p_0(n) \Lambda - \sum_{k=1}^{\infty} a_k \nabla[k-1] p_0(n) \Lambda^{-k+1} \\ &= \sum_{k=1}^{\infty} \delta_n a_k \Lambda^{-k+1} - [p_0(n), L] = \varepsilon^{-1}[B_n, L] - [p_0(n), L] \\ &= \varepsilon^{-1}[\overline{C_n}, \bar{L}]. \end{aligned}$$

A similar proof shows that $\overline{\bar{\delta}_n L} = \varepsilon^{-1}[\overline{C_n}, L]$. □

It is automatic that the derivations $\bar{\delta}_m$ and $\bar{\delta}_n$ commute, since their conjugates do. To see that δ_m commutes with $\bar{\delta}_n$, we use the Zakharov-Shabat equation

$$(12) \quad \delta_m \bar{B}_n - \bar{\delta}_n B_m = \varepsilon^{-1}[B_m, \bar{B}_n].$$

This is proved by combining the equations

$$\delta_m \bar{B}_n = (\delta_m \bar{L}^n)_- = \varepsilon^{-1}[B_m, \bar{L}^n]_- = \varepsilon^{-1}[B_m, \bar{B}_n]_-,$$

and

$$\bar{\delta}_n B_m = (\bar{\delta}_n L^m)_+ = \varepsilon^{-1}[\bar{B}_n, L^m]_+ = \varepsilon^{-1}[\bar{B}_n, B_m]_+.$$

It follows from (12) that δ_m and $\bar{\delta}_n$ commute:

$$\begin{aligned} [\delta_m, \bar{\delta}_n] L &= \varepsilon^{-1} \delta_m [\bar{B}_n, L] - \varepsilon^{-1} \bar{\delta}_n [B_m, L] \\ &= \varepsilon^{-1} [\delta_m \bar{B}_n, L] + \varepsilon^{-2} [\bar{B}_n, [B_m, L]] - \varepsilon^{-1} [\bar{\delta}_n B_m, L] - \varepsilon^{-2} [B_m, [\bar{B}_n, L]] \\ &= \varepsilon^{-1} [\delta_m \bar{B}_n - \bar{\delta}_n B_m, L] + \varepsilon^{-2} [\bar{B}_n, [B_m, L]] - \varepsilon^{-2} [B_m, [\bar{B}_n, L]] = 0. \end{aligned}$$

4. THE EQUIVARIANT TODA LATTICE

Informally, a reduction of the Toda lattice is an invariant submanifold of the configuration space fixed by the involution; we formalize this as follows.

Definition 4.1. A **reduction** of the Toda lattice is a differential ideal $\mathcal{I} \subset \mathcal{A}$ invariant under conjugation and preserved by the action of the derivations δ_n and $\bar{\delta}_n$.

The simplest example of such a reduction is the **Toda chain**, defined by the constraint $L = \bar{L}$; the associated differential ideal

$$\mathcal{I} = (v - \bar{v}, a_2 - q, a_k \mid k > 2)$$

is generated by the coefficients of $L - \bar{L}$. To see that $\mathcal{I}$ is closed under the action of the derivations δ_n , it suffices to observe that the operator $L - \bar{L}$ satisfies the Lax equation $\delta_n(L - \bar{L}) = \varepsilon^{-1}[B_n, L - \bar{L}]$, and that the coefficients of $\varepsilon^{-1}[B_n, L - \bar{L}]$ are contained in the differential ideal generated by the coefficients of $L - \bar{L}$.

The constraint $L = \bar{L}$ is equivalent to the relation $\delta_n = \bar{\delta}_n$ among the Toda flows, for all n ; in particular, the Toda equation (2) becomes in this limit the equation $\delta_1^2 \log q = \nabla^2 q$.

In this paper, we study a reduction of the Toda lattice which is a deformation of the Toda chain. Let $\mathcal{A}[\nu]$ be the extension of the differential algebra $\mathcal{A}$ by a variable ν , such that $\partial\nu = 0$ and $\bar{\nu} = -\nu$, and consider families of reductions of the Toda lattice parametrized by $\mathcal{A}[\nu]$; that is, we consider differential ideals in $\mathcal{A}[\nu]$ satisfying the conditions of Definition 4.1.

Definition 4.2. The **equivariant Toda lattice** is the reduction of the Toda lattice defined over $\mathcal{A}[\nu]$ by the constraints

$$(13) \quad (\delta_1 - \bar{\delta}_1)L = \nu\partial L, \quad (\delta_1 - \bar{\delta}_1)\bar{L} = \nu\partial\bar{L}.$$

Let $\mathcal{I}_\nu$ be the differential ideal determining this reduction.

Let K be the difference operator

$$(14) \quad K = B_1 - C_1 = \Lambda + v + q\Lambda^{-1}.$$

Substituting the Lax equations into the constraints (13), we obtain an equivalent formulation of the equivariant Toda lattice: it is characterized by the pair of equations

$$(15) \quad \varepsilon^{-1}[K, L] = \nu\partial L, \quad \varepsilon^{-1}[K, \bar{L}] = \nu\partial\bar{L}.$$

In other words, the differential ideal $\mathcal{I}_\nu$ defining the equivariant Toda lattice is generated by the coefficients of these equations.

The following theorem collects the main properties of the differential ideal $\mathcal{I}_\nu$.

Theorem 4.1. *Let $P : \mathcal{A} \rightarrow \mathcal{A}$ be the infinite-order differential operator*

$$\begin{aligned} P &= \frac{\partial}{\nabla} = \sum_{g=0}^{\infty} \frac{\varepsilon^{2g}(2^{1-2g} - 1)B_{2g}}{(2g)!} \partial^{2g} \\ &= 1 - \frac{1}{24}\varepsilon^2 \partial^2 + O(\varepsilon^4). \end{aligned}$$

The differential ideal $\mathcal{I}_\nu$ defining the equivariant Toda lattice equals $(y, \partial z_k \mid k > 0)$, where

$$y = q\nabla(v - \bar{v}) - \nu\partial q,$$

$$z_k = p_{-1}(k) - qp_1(k) - \nu Pp_0(k).$$

The differential algebra $\mathcal{A}[\nu]/\mathcal{I}_\nu$ is isomorphic to

$$\tilde{\mathcal{A}} = \mathbb{Q}_{\varepsilon, \nu} \{q, v, z_k \mid k > 0\} / (q - \bar{q}, y, \partial z_k \mid k > 0).$$

The Toda flows δ_n and $\bar{\delta}_n$ map z_k to $\mathcal{I}_\nu$.

Proof. Let $\tilde{\mathcal{I}}_\nu$ be the differential ideal $(y, \partial z_k \mid k > 0)$. Define elements $f_k(n), g_k(n) \in \mathcal{A}$ by the formulas

$$\varepsilon^{-1}[K, L^n] - \nu \partial L^n = \sum_{k=-\infty}^{\infty} f_k(n) \Lambda^{-k}, \quad \varepsilon^{-1}[K, \bar{L}^n] - \nu \partial \bar{L}^n = \sum_{k=-\infty}^{\infty} g_k(n) \Lambda^k.$$

The differential ideal $\mathcal{I}_\nu$ is generated by the coefficients $f_k = f_k(1)$ and $g_k = g_k(1)$. The formulas

$$\begin{aligned} \varepsilon^{-1}[K, L^n] - \nu \partial L^n &= \sum_{i=1}^n L^{i-1} (\varepsilon^{-1}[K, L] - \nu \partial L) L^{n-i}, \\ \varepsilon^{-1}[K, \bar{L}^n] - \nu \partial \bar{L}^n &= \sum_{i=1}^n \bar{L}^{i-1} (\varepsilon^{-1}[K, \bar{L}] - \nu \partial \bar{L}) \bar{L}^{n-i}, \end{aligned}$$

show that the coefficients $f_k(n)$ and $g_k(n)$ lie in $\mathcal{I}_\nu$; hence $\partial z_n = \mathbf{P}f_0(n)$ and $\partial \bar{z}_n = \mathbf{P}g_0(n)$ do as well, showing that $\tilde{\mathcal{I}}_\nu \subset \mathcal{I}_\nu$. We wish to prove the equality of these two differential ideals.

To do this, we show, by induction on k , that the coefficients f_k lie in $\tilde{\mathcal{I}}_\nu$. We have $f_k = 0$ for $k < 0$. If $f_j \in \tilde{\mathcal{I}}_\nu$ for $j < k$, we see that

$$\begin{aligned} \nabla z_{k+1} &= \text{res}(\varepsilon^{-1}[K, L^{k+1}] - \nu \partial L^{k+1}) = \sum_{i=0}^k \text{res}(L^i (\varepsilon^{-1}[K, L] - \nu \partial L) L^{k-i}) \\ &\equiv [k+1]f_k \pmod{\tilde{\mathcal{I}}_\nu}, \end{aligned}$$

hence $f_k \in \tilde{\mathcal{I}}_\nu$. A similar induction shows that $g_k \in \tilde{\mathcal{I}}_\nu$; this induction starts with the fact that $g_{-1} = y$ lies in $\tilde{\mathcal{I}}_\nu$.

Since $z_k - [k]a_{k+1} \in (q, v, a_2, \dots, a_k)$, we see that the differential algebra $\mathcal{A}/\mathcal{I}_\nu$ is isomorphic to

$$\tilde{\mathcal{A}} = \mathbb{Q}_{\varepsilon, \nu} \{q, v, z_k \mid k > 0\} / (q - \bar{q}, y, \partial z_k \mid k > 0).$$

It remains to prove that $\delta_n z_k$ and $\bar{\delta}_n z_k$ lie in $\mathcal{I}_\nu$. By the Zakharov-Shabat equations (11) and (12), we see that

$$\begin{aligned} \delta_n K &= \delta_n (B_1 - C_1) = (\delta_1 - \bar{\delta}_1) B_n + \varepsilon^{-1}[B_n, B_1 - C_1] \\ &= \nu \partial B_n + \varepsilon^{-1}[B_n, K]. \end{aligned}$$

It follows that

$$\begin{aligned}
\nabla \delta_n z_k &= \delta_n f_0(k) = \delta_n \operatorname{res}(\varepsilon^{-1}[K, L^k] - \nu \partial L^k) \\
&= \operatorname{res}(\varepsilon^{-1}[\delta_n K, L^k] + \varepsilon^{-1}[K, \delta_n L^k] - \nu \partial \delta_n L^k) \\
&= \varepsilon^{-1} \operatorname{res}([\nu \partial B_n + \varepsilon^{-1}[B_n, K], L^k] + \varepsilon^{-1}[K, [B_n, L^k]] - \nu \partial [B_n, L^k]) \\
&= \varepsilon^{-1} \operatorname{res}([B_n, \varepsilon^{-1}[K, L^k] - \nu \partial L^k]) \\
&= \nabla \sum_{j=1}^n [j] (p_j(n) f_j(k)).
\end{aligned}$$

The extension of α to a homomorphism from $\mathcal{A}[\nu]$ to $\mathbb{Q}_{\varepsilon, \nu}$ continues to satisfy (10). It follows that $\alpha(\delta_n z_k) = 0$, hence we obtain an explicit equation for $\delta_n z_k$:

$$\delta_n z_k = \sum_{j=1}^n [j] (p_j(n) f_j(k)) \in \mathcal{I}_\nu.$$

The proof that $\bar{\delta}_n z_k \in \mathcal{I}_\nu$ follows along the same lines. $\square$

Let us calculate explicitly the coefficients a_2 and a_3 of the Lax operator L as elements of $\tilde{\mathcal{A}}$. Applying $\operatorname{res} : \Phi_-(\mathcal{A}, q) \rightarrow \mathcal{A}$ to the equation $\varepsilon^{-1}[K, L] = \nu \partial L$, we see that

$$a_2 = q + \nu P v + z_1.$$

Taking the coefficient of Λ^{-1} in the equation $\varepsilon^{-1}[K, L] = \nu \partial L$, we see that

$$\nabla a_3 + (a_2 - q) \nabla v = \nu \partial a_2.$$

Lemma 4.2. $\nabla f P g = \frac{1}{2} \nabla (f[2] P g) - \frac{1}{2} [2] (f \partial g)$

Proof. We have

$$\nabla f P g = \varepsilon^{-1} \mathbf{E}^{1/2} \left(f \mathbf{E}^{-1/2} P g \right) - \varepsilon^{-1} \mathbf{E}^{-1/2} \left(f \mathbf{E}^{1/2} P g \right).$$

The result follows, since $\mathbf{E}^{\pm 1/2} P = \frac{1}{2} [2] P \pm \frac{1}{2} \varepsilon \partial$. $\square$

By this lemma,

$$(a_2 - q) \nabla v = \nu \nabla v P v + z_1 \nabla v = \nu \nabla \left(\frac{1}{2} v [2] P v - \frac{1}{4} P [2] v^2 \right) + z_1 \nabla v.$$

It follows that

$$a_3 = \nu \left(P \left(\frac{1}{4} [2] v^2 + q \right) - \frac{1}{2} v [2] P v \right) + \nu^2 P v - z_1 v + \frac{1}{2} z_2.$$

This method of calculating the coefficients a_k becomes cumbersome for larger values of k : instead, it is better to use the recursion

$$(16) \quad p_{-1}(n) = q p_1(n) + \nu P p_0(n) + z_n$$

which is a consequence of Theorem 4.1.

Let Ψ be the algebra of difference operators

$$\Psi = \{ A \in \Phi_-(\tilde{\mathcal{A}}, q) \mid \varepsilon^{-1}[K, A] = \nu \partial A \}.$$

Let $\mathbb{L} \in \Psi$ be the Lax operator defined by the recursion

$$(17) \quad p_{-1}(n) = qp_1(n) + \nu p_0(n).$$

This Lax operator plays a special role in the theory: the following lemma shows that the algebra Ψ may be identified with the commutative algebra $\tilde{\mathcal{A}}_0((\mathbb{L}))$, where

$$\tilde{\mathcal{A}}_0 = \mathbb{Q}_{\varepsilon, \nu}[z_k, \bar{z}_k \mid k > 0]$$

is the kernel of the derivation $\partial : \tilde{\mathcal{A}} \rightarrow \tilde{\mathcal{A}}$.

Lemma 4.3. *The homomorphism $\tilde{\alpha} : \tilde{\mathcal{A}} \rightarrow \tilde{\mathcal{A}}_0$ which sends the generators $\partial^n q$, $\partial^n v$ and $\partial^n \bar{v}$ of $\tilde{\mathcal{A}}$ to 0 induces an isomorphism between Ψ and $\tilde{\mathcal{A}}_0((\Lambda^{-1}))$.*

Proof. Since $\tilde{\alpha}(\mathbb{L}) = \Lambda$, the map $\tilde{\alpha} : \Psi \rightarrow \tilde{\mathcal{A}}_0((\Lambda^{-1}))$ is surjective. Suppose that $A \in \Psi$ lies in the kernel of $\tilde{\alpha}$, and let k be the smallest integer such that the coefficient $x \in \tilde{\mathcal{A}}$ of Λ^{-k} in A is nonzero. We have

$$\varepsilon^{-1}[K, A] - \nu \partial A = \nabla x \Lambda^{1-k} + O(\Lambda^{-k}),$$

hence $x \in \tilde{\mathcal{A}}_0$. In this way, we see that $\tilde{\alpha} : \Psi \rightarrow \tilde{\mathcal{A}}_0((\Lambda^{-1}))$ is injective. $\square$

Theorem 4.4. *The evolutionary derivation $e = \partial_v + \partial_{\bar{v}}$ of $\tilde{\mathcal{A}}$ preserves Ψ , and $e(L)$ satisfies the formula*

$$\left(L - \nu + \sum_{k=1}^{\infty} z_k L^{-k} \right) e(L) = L.$$

Proof. If $A \in \Psi$, we have

$$\varepsilon^{-1}[K, e(A)] - \nu \partial e(A) = e(\varepsilon^{-1}[K, A] - \nu \partial A) - [e(K), A] = 0,$$

since $e(K) = 1$. This shows that e preserves Ψ .

For $n > 0$, we have by (17) that $\tilde{\alpha}(e(p_{-1}(n))) = \nu \tilde{\alpha}(e(p_0(n)))$, or equivalently,

$$\oint (\mathbb{L} - \nu) e(\mathbb{L}^n) \frac{d\mathbb{L}}{\mathbb{L}} = 0.$$

Since Ψ is a commutative algebra, $e(\mathbb{L})$ commutes with $\mathbb{L}$, hence $e(\mathbb{L}^n) = n\mathbb{L}^{n-1}e(\mathbb{L})$, and

$$\oint (\mathbb{L} - \nu) \mathbb{L}^{n-2} e(\mathbb{L}) d\mathbb{L} = 0, \quad n > 0.$$

This shows that the coefficient of $\mathbb{L}^{-k}$ in $(\mathbb{L} - \nu)e(\mathbb{L})$ vanishes, hence $(\mathbb{L} - \nu)e(\mathbb{L}) = \mathbb{L}$.

Since $L/e(L)$ lies in Ψ , there is an expansion

$$\frac{L}{e(L)} = L - \nu + \frac{1}{2\pi i} \sum_{n=0}^{\infty} L^{-n-1} \oint L^n \frac{dL}{e(L)}.$$

(The constant term is determined by the fact that $e(L) = 1 + \nu\Lambda^{-1} + O(\Lambda^{-2})$.) We have

$$\frac{dL}{e(L)} = \frac{d\mathbb{L}}{e(\mathbb{L})} = (\mathbb{L} - \nu) \frac{d\mathbb{L}}{\mathbb{L}},$$

hence

$$\frac{1}{2\pi i} \oint L^n \frac{dL}{e(L)} = \frac{1}{2\pi i} \oint L^n (\mathbb{L} - \nu) \frac{d\mathbb{L}}{\mathbb{L}}.$$

It follows from the recursion (16) that

$$z_n = \tilde{\alpha}(p_{-1}(n)) - \nu \tilde{\alpha}(p_0(n)) = \frac{1}{2\pi i} \oint L^n (\mathbb{L} - \nu) \frac{d\mathbb{L}}{\mathbb{L}},$$

and the theorem follows. $\square$

5. THE DISPERSIONLESS LIMIT OF THE EQUIVARIANT TODA LATTICE

In this section, we consider the dispersionless limit of the equivariant Toda lattice, in which $\varepsilon \rightarrow 0$; we only consider the case in which the constants of motion z_k are set to 0. If $A \in \Phi_{\pm}(\tilde{\mathcal{A}}, q)$, we write

$$A_0 = \lim_{\varepsilon \rightarrow 0} A.$$

In the dispersionless limit, the algebra $\Phi_{-}(\tilde{\mathcal{A}}, q)$ degenerates to the commutative algebra $\tilde{\mathcal{A}}((\Lambda^{-1}))$, and the leading order in the commutator is the Poisson bracket

$$\{A_0, B_0\} = \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [A, B] = (\Lambda \partial_{\Lambda} A_0) \partial B_0 - \partial A_0 (\Lambda \partial_{\Lambda} B_0).$$

It is not hard to write down explicit formulas for the Lax operator $\mathbb{L}$ of the equivariant Toda lattice and its conjugate $\bar{\mathbb{L}}$ in the dispersionless limit.

Theorem 5.1. *We have*

$$\mathbb{L}_0 = K_0 + \nu \sum_{n=0}^{\infty} \left(\frac{\nu}{\Lambda} \right)^n \sum_{k=0}^n (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \frac{\log(K_0/\Lambda)^{n-k+1}}{(n-k+1)!} \left(\frac{K_0}{\Lambda} \right)^{-n} \in \mathbb{Q}[\nu, q, v]((\Lambda^{-1}))$$

and

$$\bar{\mathbb{L}}_0 = K_0 - \nu \sum_{n=0}^{\infty} \left(\frac{-\nu \Lambda}{q} \right)^n \sum_{k=0}^n (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \frac{\log(\Lambda K_0)^{n-k+1}}{(n-k+1)!} \left(\frac{\Lambda K_0}{q} \right)^{-n} \in \mathbb{Q}[\nu, q, \bar{v}]((\Lambda^{-1})).$$

Proof. Denote by L_0 the expression which we wish to prove equals $\mathbb{L}_0$. It is clear that $\beta(L) = \Lambda$, hence it suffices to prove the equation

$$\{K_0, L_0\} = \nu \partial K_0,$$

which is the dispersionless limit of the equation $\varepsilon^{-1} [K, \mathbb{L}] = \nu \partial \mathbb{L}$.

Since $\{K_0, \log(K_0/\Lambda)\} = \partial K_0$, we have

$$\{K_0, L_0\} = \nu \partial K_0 \sum_{n=0}^{\infty} \left(\frac{\nu}{\Lambda} \right)^n \sum_{k=0}^n (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \frac{\log(K_0/\Lambda)^{n-k}}{(n-k)!} \left(\frac{K_0}{\Lambda} \right)^{-n}.$$

Since $\partial \log(K_0/\Lambda) = K_0^{-1} \partial K_0$, we have

$$\partial L_0 = \partial K_0 \left(1 + \sum_{n=0}^{\infty} \left(\frac{\nu}{\Lambda} \right)^n \sum_{k=0}^n (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \left(\frac{\log(K_0/\Lambda)^{n-k}}{(n-k)!} - \frac{n \log(K_0/\Lambda)^{n-k+1}}{(n-k+1)!} \right) \left(\frac{K_0}{\Lambda} \right)^{-n-1} \right).$$

The equation $\{K_0, L_0\} = \nu \partial L_0$ follows from the recursion $\begin{bmatrix} n \\ k \end{bmatrix} = (n-1) \begin{bmatrix} n-1 \\ k \end{bmatrix} + \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}$.

Define $e^{\circ j}(L_0)$ by induction: $e^{\circ 0}(L_0) = L_0$ and $e^{\circ(j+1)}(L_0) = e(e^{\circ j}(L_0))$. Then

$$e^{\circ j}(L_0) = \delta_{j,0} K_0 + \nu^{1-j} \sum_{n=0}^{\infty} \left(\frac{\nu}{\Lambda}\right)^n \sum_{k=0}^{n-j+1} (-1)^{n-k-j} \begin{bmatrix} n \\ k \end{bmatrix} \frac{\log(K_0/\Lambda)^{n-k-j+1}}{(n-k-j+1)!} \left(\frac{K_0}{\Lambda}\right)^{-n}.$$

This formula is proved by induction on j , using the formulas $e(K_0) = 1$ and $e(\log(K_0/\Lambda)) = K_0^{-1}$.

There is an embedding of the differential algebra $\tilde{\mathcal{A}}$ in the differential algebra

$$\tilde{\mathcal{A}}\{u\}/(\partial q - q\partial u) \cong \mathbb{Q}_{\varepsilon, \nu}[q]\{u, v\},$$

given by mapping $\bar{v}$ to $v - \nu Pu$. In the dispersionless limit, this embedding maps $\bar{v}$ to $v - \nu u$.

We will prove the formula for $\bar{\mathbb{L}}_0$ by working with Laurent series in this larger algebra.

The Laurent series $\bar{\mathbb{L}}_0$ is obtained from $\mathbb{L}_0$ by replacing v by $v - \nu u$, Λ by q/Λ , and ν by $-\nu$. Let $\tilde{\mathbb{L}}_0$ be the result of substituting $v - \nu u$ for v in $\mathbb{L}_0$; it is given by the formula

$$\begin{aligned} \tilde{\mathbb{L}}_0 &= \sum_{j=0}^{\infty} \frac{(-\nu u)^j}{j!} e^{\circ j}(L) \\ &= K_0 + \nu \sum_{n=0}^{\infty} \left(\frac{\nu}{\Lambda}\right)^n \sum_{j=0}^{n+1} \sum_{k=0}^{n-j+1} (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \frac{u^j \log(K_0/\Lambda)^{n-k-j+1}}{j!(n-k-j+1)!} \left(\frac{K_0}{\Lambda}\right)^{-n} \\ &= K_0 + \nu \sum_{n=0}^{\infty} \left(\frac{\nu}{\Lambda}\right)^n \sum_{k=0}^{n+1} (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \frac{\log(qK_0/\Lambda)^{n-k+1}}{(n-k+1)!} \left(\frac{K_0}{\Lambda}\right)^{-n}. \end{aligned}$$

We obtain $\bar{\mathbb{L}}_0$ on substituting $-\nu$ for ν and Λ for q/Λ . □

Define generating functions $\pi_k(z) \in \mathbb{Q}[\nu, q, v]((z))$ by the formula

$$\mathbb{L}_0(z) = \sum_{n=-\infty}^{\infty} \frac{z^n}{[n]!} \mathbb{L}_0^n = \sum_{k=-\infty}^{\infty} \pi_k(z) \Lambda^k,$$

where $[n]!$ is the rational function

$$[n]! = \frac{\Gamma(\nu z + n + 1)}{\Gamma(\nu z + 1)} = (1 + z\nu)(2 + z\nu) \dots (n + z\nu).$$

The important cases for us will be

$$\pi_{-1}(z) = \nu + \lim_{\varepsilon \rightarrow 0} \sum_{n=1}^{\infty} \frac{z^n p_k(n)}{[n]!}, \quad \pi_0(z) = 1 + \lim_{\varepsilon \rightarrow 0} \sum_{n=1}^{\infty} \frac{z^n p_0(n)}{[n]!}, \quad \pi_1(z) = \lim_{\varepsilon \rightarrow 0} \sum_{n=1}^{\infty} \frac{z^n p_1(n)}{[n]!}.$$

It follows from the recursion (17) for the coefficients of $\mathbb{L}$ that

$$(18) \quad \pi_{-1}(z) = q\pi_1(z) + \nu\pi_0(z).$$

Lemma 5.2. $e(\pi_k(z)) = z\pi_k(z)$ and $\partial_q \pi_k(z) = z\pi_{k+1}(z)$.

Proof. We have

$$\begin{aligned} (\mathbb{L}_0 - \nu)e(\mathbb{L}_0(z)) &= \sum_{n=-\infty}^{\infty} \frac{nz^n}{[n]!} \mathbb{L}_0^{n-1}(\mathbb{L}_0 - \nu)e(\mathbb{L}_0) = \sum_{n=-\infty}^{\infty} \frac{nz^n}{[n]!} \mathbb{L}_0^n \\ &= \sum_{n=-\infty}^{\infty} \frac{(\nu z + n)z^n}{[n]!} \mathbb{L}_0^n - \sum_{n=-\infty}^{\infty} \frac{\nu z^{n+1}}{[n]!} \mathbb{L}_0^n = z(\mathbb{L}_0 - \nu)\mathbb{L}_0(z). \end{aligned}$$

The equation for $e(\pi_k(z))$ follows on taking the coefficient of Λ^k . Since $\Lambda\partial_q K_0 = e(K_0)$, it follows from Theorem 5.1 that $\Lambda\partial_q \mathbb{L}_0(z) = e(\mathbb{L}_0(z))$. Taking the coefficient of Λ^{k+1} , we obtain the formula for $\partial_q \pi_k(z)$. $\square$

We may now prove the formulas (5a) and (5b) relating the dispersionless limit of the equivariant Toda lattice to the equivariant genus 0 Gromov-Witten potential of $\mathbb{CP}^1$.

In the genus 0 limit, the functions $q = \exp(u)$ and v on the large phase space become $\exp(\partial^2 \mathcal{F}_0)$ and $\partial\partial_0 \mathcal{F}_0$, and $\bar{v}$ becomes $v - \nu u$. The proof of Theorem 4.2 of [5] extends to the equivariant case, and shows that

$$(19) \quad \partial^n u = s_n + O(|s|^2 + |t|^2), \quad \partial^n v = \delta_{1,n} + t_n + O(|s|^2 + |t|^2).$$

Hence, we may identify the large phase space with the space of formal jets in an affine space with coordinates u and v .

The following lemma shows that the vector field e lifts to the puncture vector field on the large phase space.

Lemma 5.3. *The puncture vector field*

$$e = \partial - \sum_{n=0}^{\infty} \left(s_{n+1} \frac{\partial}{\partial s_n} + t_{n+1} \frac{\partial}{\partial t_n} \right)$$

on the large phase space acts on elements of $\mathbb{Q}[\nu, q, v]$ by the derivation ∂_v .

Proof. Observe that the puncture vector field e commutes with $\partial 1$; this reflects the fact that $\mathbb{CP}^1$ is one-dimensional. The puncture (or string) equation says that

$$(20) \quad e(\mathcal{F}_0) = s_0 t_0 + \frac{1}{2} \nu t_0^2.$$

Applying the differential operators $\partial_0 \partial$ and ∂^2 to this equation, we see that $e(v) = 1$ and $e(u) = 0$. $\square$

Theorem 5.4. *The dispersionless limits of (5a) and (5b) hold.*

Proof. We will concentrate on the proof of (5a). The proof of the dispersionless limit of (5b) is the same, up to conjugation.

Let $\partial(z)$ be the generating function for vector fields

$$\partial(z) = \sum_{k=0}^{\infty} z^k \partial_k.$$

We must prove that

$$\partial(z)v = \sum_{n=1}^{\infty} \frac{z^{n-1} \partial p_0(n)}{[n]!}, \quad \partial(z)u = \sum_{n=1}^{\infty} \frac{z^{n-1} \partial p_{-1}(n)}{[n]!}.$$

In terms of the generating functions

$$x = 1 + z \partial(z) \partial \mathcal{F}_0 - g_0(z), \quad y = \nu + z \partial(z) \partial_0 \mathcal{F}_0 - g_{-1}(z),$$

we wish to prove that $\partial x(z) = \partial y(z) = 0$. We will actually prove the stronger result, that $x(z) = y(z) = 0$: in other words, that

$$\sum_{k=0}^{\infty} z^k \partial_k \partial \mathcal{F}_0 = \sum_{n=1}^{\infty} \frac{z^{n-1} p_0(n)}{[n]!}, \quad \sum_{k=0}^{\infty} z^k \partial_k \partial_0 \mathcal{F}_0 = \sum_{n=1}^{\infty} \frac{z^{n-1} p_{-1}(n)}{[n]!}.$$

A theorem of Dijkgraaf and Witten [2] establishes that the Toda equation (2) holds in the dispersionless limit:

$$\partial_0^2 \mathcal{F}_0 = q + \nu v.$$

Combining the topological recursion relations for equivariant Gromov-Witten invariants in genus 0 with Lemma 5.2, we see that

$$\begin{aligned} \partial x(z) &= z(x(z) \partial(v - \nu u) + y(z) \partial u) \\ \partial y(z) &= z(x(z) \partial q + y(z) \partial v). \end{aligned}$$

On the other hand, the string equation shows that $e(x(z)) = zx(z)$ and $e(y(z)) = zy(z)$.

Now apply the following principle (Proposition 4.1 of [5]):

A function f on the large phase space such that ∂f and $e(f)$ lie in $\mathbb{Q}[\nu]$ itself lies in $\mathbb{Q}[\nu]$.

Arguing by induction, we see that the coefficients of z^k in $x(z)$ and $y(z)$ lie in $\mathbb{Q}[\nu]$; in other words, $x(z), y(z) \in \mathbb{Q}[[\nu, z]]$. (In particular, we see that $\partial_k \partial \mathcal{F}_0$ and $\partial_k \partial_0 \mathcal{F}_0$ lie in $\mathbb{Q}[\nu, q, v]$ for all $k \geq 0$.)

The proof is finished by observing that, by the divisor equation for Gromov-Witten invariants, the limits $\lim_{q \rightarrow 0} x(z)$ and $\lim_{q \rightarrow 0} y(z)$ are integrals over the degree 0 moduli space $\overline{\mathcal{M}}_{0,2}(\mathbb{CP}^1, 0)$; however, this moduli space is empty, hence $x(z) = y(z) = 0$. $\square$

As mentioned in the introduction, the analogue of Theorem 5.4 is now known to hold in all genera (Okounkov and Pandharipande [7]).

REFERENCES

- [1] G. Carlet, B. Dubrovin and Y. Zhang, to appear.
- [2] R. Dijkgraaf and E. Witten, *Mean field theory, topological field theory, and multi-matrix models*. Nucl. Phys. **B 342** (1990), 486–522.
- [3] T. Eguchi and S.-K. Yang, *The topological $\mathbb{CP}^1$ model and the large- N matrix integral*. Modern Phys. Lett. **A 9** (1994), 2893–2902. <hep-th/9407134>

- [4] T. Eguchi, K. Hori and S.-K. Yang, *Topological σ -models and large- N matrix integral*. Internat. J. Modern Phys. **A 10** (1995), 4203–4224. <hep-th/9503017>
- [5] E. Getzler, *The Toda conjecture*. In “Symplectic geometry and mirror symmetry (KIAS, Seoul, 2000),” eds. K. Fukaya et al., World Scientific, Singapore, 2001, pp. 51–79. <math/0108108>
- [6] B. A. Kupershmidt, “Discrete Lax equations and differential-difference calculus.” Astérisque **123** (1985).
- [7] A. Okounkov and R. Pandharipande, *The equivariant Gromov-Witten theory of $\mathbf{P}^1$* . <math/0207233>
- [8] R. Pandharipande, *The Toda equations and the Gromov-Witten theory of the Riemann sphere*. Lett. Math. Phys. **53** (2000), 59–74. <math/9912166>
- [9] R. Pandharipande, private communication (2000).
- [10] K. Takasaki and T. Takebe, *Quasi-classical limit of Toda hierarchy and W -infinity symmetries*. Lett. Math. Phys. **28** (1993), 165–176.
- [11] K. Ueno and K. Takasaki, *Toda lattice hierarchy*. In “Group representations and systems of differential equations (Tokyo, 1982).” Adv. Stud. Pure Math., vol. 4, North-Holland, Amsterdam, 1984, pp. 1–95.

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, ILLINOIS